

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification 5 :</b> <b>C07D 307/93</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 93/06095</b> <b>(43) International Publication Date:</b> 1 April 1993 (01.04.93)
<b>(21) International Application Number:</b> PCT/US92/07768 <b>(22) International Filing Date:</b> 14 September 1992 (14.09.92)  <b>(30) Priority data:</b> 764,181 23 September 1991 (23.09.91) US  <b>(71) Applicant:</b> IOWA STATE UNIVERSITY RESEARCH FOUNDATION, INC. [US/US]; 214 O&L, Ames, IA 50011-3020 (US).  <b>(72) Inventors:</b> LAROCK, Richard, C. ; Rural Route 4, Ames, IA 50010 (US). LEE, Nam, Ho ; 2022A Orchard Street, Urbana, IL 61801 (US).  <b>(74) Agent:</b> HAMRE, Curtis, B.; Merchant, Gould, Smith, Edell, Welter & Schmidt, 3100 Norwest Center, 90 South Seventh Street, Minneapolis, MN 55402 (US).		<b>(81) Designated States:</b> JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> FREE RADICAL-CATALYZED SYNTHESIS OF BENZOPROSTACYCLINS  <b>(57) Abstract</b>  A method is provided for preparing benzoprostacyclins by the free-radical-catalyzed tandem alkene insertion into a 1,4-bi-soxy-substituted cyclopent-2-ene intermediate.		

BEST AVAILABLE COPY

**FOR THE PURPOSES OF INFORMATION ONLY**


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MN	Mongolia
AU	Australia	FR	France	MR	Mauritania
BB	Barbados	GA	Gabon	MW	Malawi
BE	Belgium	GB	United Kingdom	NL	Netherlands
BF	Burkina Faso	GN	Guinea	NO	Norway
BG	Bulgaria	GR	Greece	NZ	New Zealand
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	PT	Portugal
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	RU	Russian Federation
CG	Congo	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovak Republic
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CS	Czechoslovakia	LU	Luxembourg	SU	Soviet Union
CZ	Czech Republic	MC	Monaco	TD	Chad
DE	Germany	MG	Madagascar	TG	Togo
DK	Denmark	MI	Mali	UA	Ukraine
ES	Spain			US	United States of America


**BEST AVAILABLE COPY**

WHAT IS CLAIMED IS:


1. A method for preparing benzoprostacyclins comprising reacting a compound of the general formula (III):



- wherein  $R^1$  is a pharmaceutically-acceptable cation, H or  $(C_1-C_{12})$ alkyl; A is  $-CH_2-$ ,  $-O-CH_2-$ ,  $-CH_2-CH_2-$  or  $-CH=CH-$ ; and  $R^2$  is H,  $(C_1-C_{12})$ alkyl,  $(C_1-C_{10})$ acyl or  $(C_7-C_{13})$ aroyl with a compound of a general formula (IV):
- 15



- 20 wherein  $R^5$  is  $(C_1-C_5)$ alkyl; B is  $-(CH_2)_n-Z$  wherein n is 0-4 and Z is  $-CH_2CH_2-$ ,  $-CH=CH-$  or  $-C\equiv C-$ ;  $R^4$  is H, F,  $CH_3$  or  $CH_2CH_3$ ; and  $R^7$  is  $((C_1-C_4)$ alkyl) $_3$ Sn or (phenyl) $_3$ Sn wherein the reaction is carried out in the presence of a catalytic amount of a free radical initiator to yield a compound of the formula (II):
- 25




- wherein  $R^8$  and  $R^9$  taken together are keto, and  $R^1$ , A,  $R^2$ ,  $R^4$ , B and  $R^5$  are as defined above.


SUBSTITUTE SHEET

BEST AVAILABLE COPY

2. The method of claim 1, further comprising reducing the C<sub>15</sub>-keto group of compound II with a reducing agent to yield a compound of formula II wherein R<sup>8</sup> is H and R<sup>9</sup> is OH.
3. The method of claim 1 wherein, in compound III, R<sup>1</sup> is (C<sub>1</sub>-C<sub>12</sub>)alkyl and R<sup>2</sup> is H.
4. The method of claim 1 wherein the mole ratio of III:IV is about 1:1.25-20.
5. A method for preparing benzoprostacyclins comprising reacting a compound of the general formula (III):



wherein R<sup>1</sup> is a pharmaceutically acceptable cation, H or (C<sub>1</sub>-C<sub>12</sub>)alkyl; A is -CH<sub>2</sub>-, -O-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>- or -CH=CH-; and R<sup>2</sup> is H, (C<sub>1</sub>-C<sub>12</sub>)alkyl, (C<sub>1</sub>-C<sub>10</sub>)acyl or (C<sub>7</sub>-C<sub>13</sub>)aroyl with a compound of the general formula (V):




wherein R<sup>5</sup> is (C<sub>1</sub>-C<sub>5</sub>)alkyl, B is -(CH<sub>2</sub>)<sub>n</sub>-Z wherein n is 0-4 and Z is -CH<sub>2</sub>CH<sub>2</sub>-, -CH=CH- or -C≡C-; R<sup>4</sup> is H, F, CH<sub>3</sub> or CH<sub>2</sub>CH<sub>3</sub>; R<sup>7</sup> is ((C<sub>1</sub>-C<sub>4</sub>)alkyl)<sub>3</sub>Sn or (phenyl)<sub>3</sub>-Sn, R<sup>8</sup> is (C<sub>1</sub>-C<sub>12</sub>)alkyl or H; and R<sup>3</sup> is H, (C<sub>1</sub>-C<sub>12</sub>)alkyl, (C<sub>1</sub>-C<sub>10</sub>)acyl or (C<sub>7</sub>-C<sub>13</sub>)aroyl; wherein the reaction is carried out in the

BEST AVAILABLE COPY

SUBSTITUTE SHEET

presence of a catalytic amount of a free radical initiator to yield a compound of the formula II:



wherein  $R^9$  is  $OR^3$ , and  $R^1$ , A,  $R^2$ ,  $R^3$ ,  $R^4$ , B,  $R^5$  and  $R^8$  are as defined above.

- 15 6. The method of claims 1 or 5 wherein  $R^7$  is (n-butyl)<sub>3</sub>Sn.
7. The method of claim 5 wherein  $R^3$  is H in compounds II and V.
- 20 8. The method of claims 1 or 5 wherein compound II comprises (S)C<sub>15</sub>-OH.
9. The method of claim 8 wherein the C<sub>11</sub>-OR<sup>2</sup> bond is in the alpha-configuration.
- 25 10. The method of claims 1 or 5 further comprising saponifying the CO<sub>2</sub>R<sup>1</sup> moiety of compound II and neutralizing the reaction mixture to yield CO<sub>2</sub>H.
- 30 11. The method of claim 10 further comprising forming a pharmaceutically acceptable alkali metal salt, ammonium, or amine salt of the moiety CO<sub>2</sub>H.
- 35 12. The method of claims 1 or 5 wherein the free radical initiator is AIBN.

13. The method of claims 1 or 5 wherein the reaction is carried out in solution in an organic solvent.
14. The method of claim 13 wherein the reaction is carried out at about 50-150°C.
15. The method of claim 14 wherein the reaction is carried out for about 5-48 hours.
16. The method of claims 1 or 5 wherein A is  $-\text{CH}_2-$  or  $-\text{CH}_2-\text{CH}_2-$  and B is  $\text{CH}_2-\text{CH}_2-\text{CH}_2$ .
17. The method of claim 16 wherein  $\text{R}^4$  is H and  $\text{R}^5$  is  $\text{CH}_3$ .

BEST AVAILABLE COPY